


**Research Article**

## An Illustrative Case Report of Anterior Migration of Optic Disc Drusen in A Child

 Aoife Naughton\*<sup>1</sup> and Ammar Miri<sup>2</sup>

### Abstract

Optic disc drusen (ODD) are acellular calcified deposits located both intracellularly and extracellularly within the optic nerve. Their appearance typically change over the first two decades as they migrate anteriorly towards the optic nerve head, from an initial buried location. They can pose a diagnostic challenge as can simulate the appearance of optic disc swelling. Distinguishing ODD from true optic disc swelling is of paramount importance to avoid delayed diagnosis of potentially life threatening neurologic disease and on the other hand prevent invasive and unnecessary work-up for elevated intracranial pressure, if not clinically indicated. A number of imaging modalities are available to assist in the diagnosis and are discussed below.

**Keywords:** Optic disc drusen (ODD), Ophthalmology, EDI-OCT, OCT RNFL

### Introduction

Optic disc drusen (ODD) are acellular calcified deposits located both intracellularly and extracellularly within the optic nerve. Their appearance typically change over the first two decades as they migrate anteriorly towards the optic nerve head, from an initial buried location. They can pose a diagnostic challenge as can simulate the appearance of optic disc swelling. Distinguishing ODD from true optic disc swelling is of paramount importance to avoid delayed diagnosis of potentially life threatening neurologic disease and on the other hand prevent invasive and unnecessary work-up for elevated intracranial pressure, if not clinically indicated. A number of imaging modalities are available to assist in the diagnosis and are discussed below.

### Methods

We present an illustrative case of the change in appearance of ODD due to anterior migration over a 3-year period, in a young child.

### Results / Case report

A young female aged 3 years was referred by her optician to the Ophthalmology department for urgent review and evaluation of blurred disc margins. Her parents had taken her to the optician regarding a recent onset squint, which was first noticed 3 weeks previously. She had been well up until the day of her review, during which she was feeling nauseous and vomited on four occasions. She had no significant past medical history. She was born by elective caesarean section at full term, weighing 4.28kg. She was on no regular medications. There was a positive family history of strabismus.

On examination, her logMAR visual acuities were 0.175 and 0.275 in the

### Affiliation:

<sup>1</sup>Oxford Eye Hospital, John Radcliffe Hospital, Oxford, United Kingdom

<sup>2</sup>Milton Keynes University Hospital, Milton Keynes, United Kingdom

### Corresponding author:

Aoife Naughton, Oxford Eye Hospital, John Radcliffe Hospital, Headley Way, Headington, Oxford, OX3 9DU, United Kingdom.

**Citation:** : Aoife Naughton, Ammar Miri. An Illustrative Case Report of Anterior Migration of Optic Disc Drusen in A Child. Journal of Ophthalmology and Research 5 (2022): 127-130.

**Received:** September 25, 2022

**Accepted:** October 03, 2022

**Published:** October 18, 2022



important to consider optic disc drusen in the differential for papilledema, as 50 to 55% of children initially diagnosed with papilledema have optic disc drusen as their final diagnosis [3]. Features in favour of papilloedema include obscuration of peripapillary vessels caused by nerve fibre layer oedema involving the peripapillary retina, hyperemia, hemorrhages, cotton wool spots, Paton lines, and exudates around the optic disc.

ODD are commonly buried when occurring in children. They can therefore be particularly difficult to distinguish from true optic nerve swelling. Few reports have demonstrated the evolution of ODD in children over time [4-7], progressing from anomalous disc appearance to ophthalmoscopically visible optic nerve drusen. However, there is a paucity of photographic documentation of the natural history of changes in the optic disc appearance over time, particularly in the earlier stages whilst the ODD remain buried. Opportunities to document the full panorama of drusen evolution in a particular patient are limited. Our case demonstrates a marked increase in elevation of both optic discs, attributable to drusen migration, without the drusen being ophthalmoscopically visible.

Determining a definitive diagnosis based on clinical examination alone can be challenging for even the most distinguished ophthalmologists. In these cases, a variety of imaging modalities may be employed to evaluate for the presence of ODD. B-scan ultrasonography has long been the preferred method for diagnosis of ODD [8]. Drusen characteristically appear hyperechoic with posterior shadowing on ultrasonography. Leon and colleagues reported on the cost-effectiveness of performing ultrasonography prior to neuroimaging in patients with ODD, with an overall reduction in the number of other investigations undertaken [9]. However, drusen and raised intracranial pressure can coexist [10, 11] and a diagnosis of drusen does not preclude the need for investigations for papilledema. In patients with signs or symptoms suspicious for another disorder, further evaluation may be necessary even if optic disc drusen are detected. Mosajee et al found the prevalence of coexisting ODD and raised intracranial pressure in children to be 10% in their pediatric practice [12].

Mosajee et al [12] also highlighted the importance of using ultrasonography not only to detect drusen but also to measure optic nerve sheath diameter, which if large is indicative of papilloedema. Carter et al [13]. Described a method of using ultrasound to identify fluid around the optic nerve (crescent sign) as a reliable sign of optic disc swelling. A-scan mode is used to determine the optic nerve sheath width in the primary gaze. If the optic nerve sheath was found to be wide ( $\geq 3.3$  mm), a 30° test was then performed to differentiate fluid from solid thickening of the optic nerve sheath. The patient is instructed to re-fixate at least 30° away from the primary gaze

toward the ultrasound probe, at which point the optic nerve sheath was measured again. If fluid is present, the measured width of the optic nerve is expected to decrease, based on the assumption that when the eye is turned the optic nerve sheaths become stretched and the fluid is redistributed over a greater area. A reduction in width  $>10\%$  is considered positive for the presence of fluid.

In more recent years, there has been considerable interest in the use of OCT in the diagnosis of ODD. Spectral-domain OCT (SD-OCT) offers a limited depth of imaging and is considered unreliable at distinguishing between buried optic disc drusen and true optic disc edema in children [14]. The introduction of enhanced depth imaging optical coherence tomography (EDI-OCT) has improved the visualization of more deeply buried ODD [15]. EDI-OCT allows for evaluation of the entire nerve head up to the lamina cribrosa which is hypothesized to be the most posterior extent where ODD may be found. The Optic Disc Drusen Studies Consortium recommends obtaining EDI-OCT (for identification of ONHD and measurement of scleral canal size), OCT RNFL, OCT Macula, and fundus autofluorescence [16]. Autofluorescence, however, does not reliably detect buried drusen [17]. Therefore, its usefulness in identifying optic disc drusen in children, who are more likely to harbour non-calcified buried drusen, has not been conclusively determined.

## Learning points

This case illustrates the migration of optic disc drusen over a 3-year period in a child, where the margins of the discs over this time become more elevated and indistinct. EDI-OCT show a significant increase in the elevation of the optic nerve heads bilaterally, whereby it is difficult to outrule concurrent papilloedema. As discussed, ODD and papilloedema can coexist and it is therefore pertinent to take the full clinical picture into account when evaluating similar patients. The changes documented in this case were attributed to ODD migration, following full investigation, including neuroimaging and CSF sampling, as this was considered clinically indicated.

## Conflict of interest

The authors report no conflict of interest.

## Funding/Support

None to declare

## Abbreviations

ODD: optic disc drusen, ONH: optic nerve head, OCT: optical coherence tomography, SD-OCT: spectral-domain optical coherence tomography, EDI-OCT: enhanced depth imaging optical coherence tomography

## References

- Muller H. Anatomische Beiträge zur Ophthalmologie. *Albrecht von Graefes Arch Klin Ophthalmol* 4 (1858): 1–40.
- Hoover DL, Robb RM, Petersen RA. Optic disc drusen in children. *J Pediatr Ophthalmol Strabismus* 25 (1988): 191–195.
- Kovarik JJ, Doshi PN, Collinge JE, Plager DA. Outcome of pediatric patients referred for papilledema. *J AAPOS* 19 (2015): 344–348.
- Mustonen E, Alanko HI, Nieminen H. Changes in optic disc drusen. Demonstration by stereophotographs and electronic subtraction. *Acta Ophthalmol (Copenh)* 60 (1982): 3–15.
- Miller NR. Appearance of optic disc drusen in a patient with anomalous elevation of the optic disc [letter]. *Arch Ophthalmol* 104 (1986): 794–5.
- Spencer TS, Katz BJ, Weber SW, Digre KB. Progression from anomalous optic discs to visible optic disc drusen. *J Neuroophthalmol* 24 (2004): 297.
- Frisén L. Evolution of drusen of the optic nerve head over 23 years. *Acta Ophthalmol* 86 (2008): 111–112.
- Gise R, Gaier ED, Heidary G. Diagnosis and Imaging of Optic Nerve Head Drusen. *Seminars in Ophthalmology* 34 (2019): 256–263.
- Leon M, Hutchinson AK, Lenhart PD, Lambert SR. The cost-effectiveness of different strategies to evaluate optic disk drusen in children. *J AAPOS* 18 (2014): 449–452.
- Komur M, Sari A, Okuyaz C. Simultaneous papilledema and optic disc drusen in a child. *Pediatr Neurol* 46 (2012): 187–188.
- Rossiter JD, Lockwood AJ, Evans AR. Coexistence of optic disc drusen and idiopathic intracranial hypertension in a child. *Eye* 19 (2005): 234–235.
- Moosajee M, Restori M, Acheson J, Dahlmann-Noor A. The cost-effectiveness of different strategies to evaluate optic disk drusen in children Response. *J AAPOS* 19 (2015): 392-3.
- Carter SB, Pistilli M, Livingston KG, et al. The role of orbital ultrasonography in distinguishing papilledema from pseudopapilledema *Eye* 28 (2014): 1425-30.
- Kulkarni KM, Pasol J, Rosa PR, Lam BL. Differentiating mild papilledema and buried optic nerve head drusen using spectral domain optical coherence tomography. *Ophthalmology* 121 (2014): 959–963.
- Merchant KY, Su D, Park SC, Qayum S, Banik R, Liebmann JM, Ritch R. Enhanced depth imaging optical coherence tomography of optic nerve head drusen. *Ophthalmology* 120 (2013): 1409–1414.
- Malmqvist L, Bursztyn L, Costello F, et al. The Optic Disc Drusen studies consortium recommendations for diagnosis of Optic Disc Drusen using optical coherence tomography. *J Neuroophthalmol* 38 (2018): 299–307.
- Kurz-Levin MM, Landau K. A comparison of imaging techniques for diagnosing drusen of the optic nerve head. *Arch Ophthalmol* 117 (1999): 1045–1049.