





















- Hyaluronic Acid, and Two Subcomponents of Hyaluronic Acid: Synovial Fluid Scavenging Effect is Enhanced in Rheumatoid Arthritis Patients. *Arthritis Rheumatol* 31 (1988): 63-71.
28. Filippin LI, Vercelino R, Marroni NP, Xavier RM. Redox signalling and the inflammatory response in rheumatoid arthritis. *Clin Exp Immunol* 152 (2008): 415-422.
29. Venkatesha SH, Berman BM, Moudgil KD. Herbal Medicinal Products Target Defined Biochemical and Molecular Mediators of Inflammatory Autoimmune Arthritis. *Bioorgan Med Chem* 19 (2011): 21-29.
30. Xing R, Liu S, Wang L, Cai S, Yu H, Feng J, et al. The Preparation and Antioxidant Activity of Glucosamine Sulfate. *Chin J of Oceanol Limn* 27 (2009): 283-287.
31. Kokkiripati PK, Bhakshu LM, Marri S, Padmasree K, Row AT, Raghavendra AS, et al. Gum resin of *Boswellia serrata* inhibited human monocytic (THP-1) cell activation and platelet aggregation. *J Ethnopharmacol* 137 (2011): 893-901.
32. Kaur A, Nain P, Nain J. Herbal plants used in treatment of rheumatoid arthritis: a review. *Int J Pharm Pharm Sci* 4 (2012): 44-57.