


**Research Article**

## Regeneration of Medicinal Plant *Solanum virginianum* L. Via Somatic Embryogenesis from Seed Explant

Dhanashree Subhash Patil and Swaroopa Amit Patil\*

### Abstract

*Solanum virginianum* L., belonging to the family Solanaceae is an important medicinal plant, used in traditional ayurvedic medicines. The present study was aimed to develop an efficient procedure for inducing somatic embryogenesis followed by regeneration in *S. virginianum*. The seeds of *S. virginianum* were used as explant and inoculated on ½ MS (Murashige and Skoog) basal medium. Stem explant from *in vitro* germinated seeds was used for callus induction. MS medium supplemented with 1 mg L<sup>-1</sup> BAP (6- Benzylaminopurine) was used for callus induction. Callus proliferation started when initiated callus was sub-cultured on various concentrations and combinations of BAP, IAA (Indole-3-Acetic Acid), IBA (Indole-3-Butyric Acid), 2,4-D (2,4-Dichlorophenoxyacetic acid) and NAA (Naphthalene acetic acid). The highest embryogenesis took place in BAP (@ 2 mg L<sup>-1</sup>) + IAA (@ 5 mg L<sup>-1</sup>) combination. The somatic embryos were transferred onto MS basal medium for maturation. These embryos matured and were converted into complete plantlets with well-developed shoot and root system. Highest survival percentage (98±1.24%) was observed on red soil : sand (1:1) potting mixture used for acclimatization.

**Keywords:** 2,4-D, BAP, IAA, IBA, *Solanum virginianum*, somatic embryogenesis, NAA.

### Introduction

India has largest biodiversity in medicinal plants and has high knowledge of rich ancient traditional system of medicines (Ayurveda, Siddha, Unani, Amchi and local health traditions) which provides a strong base for their utilization. *Solanum virginianum* L., belongs to the family Solanaceae, is an important medicinal plant species in ayurveda and folklore medicine since time immemorial [1]. *S. virginianum* is commonly known as yellow berried night shade. *Solanum surattense* Burm. (F) and *Solanum xanthocarpum* Schrad. and Wendl. are the synonyms of *Solanum virginianum* [2]. It is widely distributed in Asia including India, China, Nepal, Pakistan, Sri Lanka, Bangladesh, Afghanistan, Thailand etc. In India it is found in tropical, subtropical, and all the four geographical regions mostly in dry places. Morphologically *S. virginianum* is a prickly diffuse bright green perennial herb which is erect or creeping. Branches are numerous and young ones are covered with dense, stellate and tomentose hairs [3].

*S. virginianum* is one of the ten constituents of “Dashmoola”. Ayurvedic physicians commonly use Dashmoola in their practices. *S. virginianum* is a chemically valuable source of alkaloids, sterols, saponins, flavonoids and their glycosides [4], containing carbohydrates, fatty acids and amino acids

#### Affiliation:

Department of Botany, Shivaji University, Kolhapur, India

#### \*Corresponding author:

Swaroopa Amit Patil, Department of Botany, Shivaji University, Kolhapur, India

**Citation:** Dhanashree Subhash Patil, Swaroopa Amit Patil. Regeneration of Medicinal Plant *Solanum virginianum* L. Via Somatic Embryogenesis from Seed Explant. Archives of Internal Medicine Research 5 (2022): 488-493.

**Received:** October 15, 2022

**Accepted:** October 27, 2022

**Published:** November 02, 2022

[5], diosgenin and beta cytosterol [6], triterpene Lupeol [7], coumarins, scopolin, scopoletin, esculin and esculetin [8], alkaloids, flavonoids and saponin and tolerable levels of heavy metals like Cu, Fe, Pb, Cd and Zn while bioactive steroidal glycoalkaloid khasianine in addition to solanine and solasomargine has been reported in *S. virginianum*. Solasodine serves as an important intermediate in the synthesis of steroidal hormone [9] and is an alternative to diosgenins as a precursor in synthesis of steroidal hormones like progesterone and cortisone [10,11]. Apigenin, another secondary metabolite from *S. virginianum* has antiallergic property while diosgenin exhibit anti-inflammatory effects [12]. Kraye and Briggs [13] reported the anti-accelerator cardiac action of solasodine and some of its derivatives. Propagation of *S. surattense*, takes place only by seeds but it faces some difficulties such as low rate of germination under normal conditions, loosing of viability on storage, the progenies produced may not be true to type due to cross pollination [14].

Globally, plant tissue culture (PTC) has been recognized for its valuable and beneficial role in the production and conservation of plant-based resources [15]. PTC can be achieved with the help of different explants such as meristem culture, anther culture, ovule culture, endosperm culture, nucellus culture, embryo culture etc. Somatic embryogenesis is the developmental process by which theoretically any somatic cell develops into a zygote like structure that finally forms a plant [16, 17] and it is significantly important for *in vitro* mass propagation, germplasm conservation and genetic improvement of higher plants.

Very limited reports are available on the *in vitro* regeneration of *S. virginianum*. Using nodal and shoot tip segments, micropropagation technique has been established in *S. virginianum* [18]. Direct shoot organogenesis from shoot tip and leaf segment was achieved by Pawar [19]. Somatic embryogenesis was successfully achieved from cotyledon and leaf explants in *S. surattense* [20]. Callus induction and shoot organogenesis from floral bud was studied [21]. Due to its over exploitation for high medicinal use, this plant is becoming endangered [22], hence there is a need for ex-situ conservation through tissue culture technique. To mitigate the natural and social problems related to *S. virginianum*, an attempt is made in the present investigation by standardizing reliable protocol for mass and rapid multiplication of this rare and endangered medicinal plant via somatic embryogenesis. The present study is focused on the effect of growth regulators that enhance somatic embryogenesis followed by shoot initiation in *S. virginianum* for mass propagation.

## Materials and Methods

Morphologically healthy fruits of *S. virginianum* were collected from Haripur, Sangli (160 50' 52.2" N 740 32' 05.8" E) (Maharashtra, India). The plantlets were maintained in the

botanical garden of Department of Botany, Shivaji University, Kolhapur. The plant was identified and authenticated by Dr. M. M. Lekhak and the herbarium sheets were prepared and submitted to Herbarium deposition section of Department of Botany, Shivaji University Kolhapur with 'SUK acronym', after which accession number SUK (DSP 01) was provided.

*In vitro* seed germination and induction of somatic embryogenesis:

All the fruits were washed with detergent Tween-20, followed by tap water twice in laboratory and surface sterilized with 0.1% HgCl<sub>2</sub> for 4 min. They were then washed thrice by sterilized distilled water for 1 min each and then treated with 70% ethanol for 30 sec. Again repeated 3 washes of sterilized distilled water for 1 min each were given. Surface sterilization was carried out in a laminar air flow. Then the fruits were blotted dry, then cut and seeds dissected out. The seeds were inoculated on to ½ strength MS medium [23] containing 3% sucrose and 0.8% Agar agar aseptically. The seeds germinated within a period of 15-20 days.

The stem from germinated seedling was used as explant and it was cultured on MS medium supplemented with BAP (1 mg L<sup>-1</sup>) for callus induction. The process of callus induction was done till the 7th week after inoculation of explant.

For induction of somatic embryogenesis the callus was transferred onto MS medium supplemented with various plant growth regulators (2,4-D, BAP, IAA, IBA and NAA) either singly or in combination (Table 1). Once the nodular embryos were formed, for further development of embryos they were transferred onto the same hormone concentration or hormone free MS medium. Experiment was repeated thrice and twenty replicates were used for each experimental set. After inoculation culture tubes were transferred to culture room under a 16/8 h (light/dark) photoperiod, temperature 25±2°C and maximum humidity (80%) was adjusted with air conditioner. The observations were recorded after nine weeks and statistical analysis was done using Microsoft excel statistical methods. The values were represented as mean ±SE.

## Results and Discussion

The important step in plant tissue culture was surface sterilization of explants. Minimum contamination and higher survival rate were achieved in 0.1% HgCl<sub>2</sub>. However, the optimum time period of exposure of seeds to HgCl<sub>2</sub> was 4 min. Surface sterilized explant was inoculated on ½ strength MS medium. Within 15 to 20 days seeds started germinating. After seed germination the elongated shoots were inoculated on different concentrations of BAP (0.5, 0.75, and 1.0 mg L<sup>-1</sup>) for multiplication. Maximum growth was achieved on 0.75mg/l BAP. BAP proved more effective for shoot regeneration than Kn for shoot multiplication of *S. virginianum* [24]. Ramar and Nandagopalan [25] confirmed

**Table 1:** Effect of different concentrations and combinations of growth regulators on embryonic callus formation in *Solanum virginianum*.

Sr. No.	Plant growth regulator (mgL <sup>-1</sup> )		Embryogenic callus formation (%)	No. of somatic embryos per tube	Callus morphology	Mean fresh wt. of callus (g)	Mean dry wt. of callus (g)
1.	MS basal medium		60	---	Whitish yellow	8.2±0.05	5.6±0.03
2.	BAP	IAA					
a.	1	3	55	4.66±0.33	Whitish green	6.0±0.04	2.5±0.30
b.	2	4	60	7.33±0.66	Whitish green	6.9±0.14	2.2±0.16
c.	2	5	90	11.66±0.33	Greenish brown	8.5±0.10	3.9±0.10
3.	BAP	IBA					
a.	1	3	55	6.33±0.33	Whitish green	5.2±0.31	2.3±0.10
b.	2	4	75	8.66±0.33	Whitish green	5.6±0.10	2.2±0.14
c.	3	5	60	6.56±0.66	Brown	7.7±0.14	3.2±0.19
4.	NAA						
a.	3	-----	65	03±0.33	Greenish brown	8.8±0.19	3.8±0.04
b.	4	-----	70	4.71±0.33	Brown	7.3±0.24	3.0±0.06
c.	5	-----	80	4.46±0.33	Brown	7.4±0.20	4.2±0.27
5.	2,4-D						
a.	3	-----	70	3.66±0.23	Greenish brown	7.2±0.10	2.9±0.18
b.	4	-----	75	5.76±0.33	Brown	7.0±0.12	2.6±0.15
c.	5	-----	80	07±0.33	Brown	6.1±0.31	2.2±0.01

**Note:** The values represent mean ± SE.

that *Solanum surattense* possess low amount of endogenous hormones hence it requires high amount of exogenous hormones for plant regeneration. The stems from germinated seedlings were inoculated on 1 mgL<sup>-1</sup>BAP which resulted in callus induction. The explants were slightly swollen and formed protuberance along whole length of the explants, which upon continuous culture gave rise to pale greenish callus within 3 to 4 weeks. The induced callus was sub-cultured on various concentrations and combinations of BAP, IAA, IBA, 2,4- D and NAA used singly or in combination (Table 1). Proliferative callus was converted into high nodular embryogenic callus in all the combinations tested. Combinations of BAP + IAA and BAP + IBA were further used for subculturing, proliferative callus was seen to rapidly convert into embryogenic callus (Figure 1a). In medium with single hormone (NAA or 2,4-D) it was seen that the proliferative callus subcultured on hormone free medium (devoid of NAA or 2,4-D) showed somatic embryogenesis (Figures 1b,1c). Thus, to initiate proliferative calli BAP, IAA, IBA, NAA and 2,4-D were used alone or in combination. For achieving somatic embryogenesis, BAP + IAA, BAP + IBA combinations were used for at the time of subculturing. All the combinations of growth regulators were more or less responsible for callus proliferation (Figures. 1b, 1c). The embryogenic calli were composed of small spherical or globular nodules. Callus formation process needed 4-6 weeks for best proliferative callus. The embryogenesis was seen in callus in the 7<sup>th</sup> week in some combinations which extended to 9<sup>th</sup> week where somatic embryogenesis was observed in

all the hormones tested. Dark conditions favored induction of somatic embryogenesis. In all the combinations of BAP+IAA studied, maximum percent response for embryogenesis (90%) was recorded in 2 mg/l BAP in combination with 5 mg/l IAA. Cytokinins, such as BAP are reported to inhibit the embryogenic potential in carrot cultures but a combination of BAP with IAA induced embryogenesis in *Podophyllum hexandrum* [26].

Embryogenic callus induction was also tested with 2,4-D and NAA alone wherein whitish callii were observed. Maximum growth was recorded on 5 mg/l for both. Ramaswamy [20] showed that leaf explants of *Solanum surattense* had maximum somatic embryogenesis and high frequency of somatic embryo induction when MS medium was supplemented with 4 mg/l NAA and 0.5 mg/l BAP. Somatic embryogenesis was reported on the medium containing NAA alone in *Solanum melongena* [27].

Less concentration of growth regulators yielded poor callus. Initial dependence of callus on growth hormone to initiate embryogenesis has been reported in a number of species but it was found necessary to withdraw the growth hormone from the medium for subsequent growth of somatic embryos. Ghatge [28] achieved initial embryogenic competence of callus in presence of 2,4-D, but found necessary to withdraw 2,4-D from the medium for further development of somatic embryos [29]. Our observations revealed that growth hormones BAP + IAA and BAP + IBA are required for the somatic embryogenesis when combinations of hormones

were used. Whereas withdrawal of growth hormones NAA and 2,4-D leads to somatic embryogenesis in case of single hormone use.

In present study, heart-shaped somatic embryos were observed which further progressed towards late-heart-shaped or torpedo-shaped embryos as they attained well-developed and mature cotyledons (Figures. 1f-k). Sharma and Millam [30] in case of *Solanum tuberosum*, observed two cotyledonary stages onto the medium supplemented with 5 µM 2,4-D. A balance in endogenous and exogenous auxin is essential for somatic embryogenesis. In carrot cultures auxin containing cultures reportedly produced more ethylene than auxin free cultures [26]. High ethylene content enhances more activity of cellulase or pectinase or both which causes the breakdown of clumps before polarity is established in proembryos for further development. Thus 2,4-D supports tissue multiplication but opposes maturity of embryos [26]. Well-developed shoots with roots were obtained after somatic embryogenesis (Figures. 1e and f), which were further dissected out and subcultured singly on MS basal medium. Cytokinins had significant effects on somatic embryogenesis and high concentration (1 mg L<sup>-1</sup>) was more effective. Similar results were obtained by Singh and Chaturvedi [12] who reported that the effects of cytokinins on somatic embryogenesis were significant and mandatory for the differentiation of somatic embryos.

In conclusion, the present study showed that the callus induction and somatic embryogenesis were strongly affected by plant growth regulators and culture conditions. The medium containing both auxin and cytokinin at any concentration is capable of inducing the formation of somatic embryogenesis but differences are based on the type and concentration of plant growth regulators and light conditions [31]. *Note: The values represent mean ± SE.*

### Hardening and acclimatization

Various potting mixtures, red soil, compost, sand coco peat (either used solely or combined) were tried for acclimatization of plants derived through somatic embryogenesis (Table 2). Well-developed plantlets were potted in plantation bags with different potting mixtures and survival percentage was calculated. All the plantation bags were kept in shade in a greenhouse in the initial days (4 weeks) to maintain humidity. Highest survival percentage (98±1.24%) was observed on red soil : sand (1:1) proportion, while lowest survival percentage (34±4.33%) was observed on Compost alone. These hardened plants were maintained in glass house for three months and then transferred to field.

For the successful multiplication of plants, micropropagation, rooting and acclimatization are important factors. In the present investigation 98 % survival rate was observed on Red soil : Sand (1:1) proportion and similar

type of results recorded by Rao [16] where tomato varieties were successfully survived on sand : soil (1:1) proportion. Large scale propagation of *S. virginianum* may be achieved by crossing the barriers like natural seed germination and seasonal restrictions.


### Conclusion

The time taken for the induction of somatic embryos was between 7 to 9 weeks. The development of somatic embryos was seen in all the cultures but with varied responses. The

**Table 2:** Effect of different potting mixtures on hardening and acclimatization of *in vitro* raised plantlets of *Solanum virginianum*.

Sr. No.	Combination	Survival percentage
1	Compost	34 ± 4.33
2	Compost : Sand (2:1)	66.9 ± 6.23
3	Red soil : Sand (1:1)	98 ± 1.24
4	Red soil	55 ± 3.98
5	Red soil : sand (2:1)	83.87 ± 2.98
6	Compost : Coco peat (1:1)	48.34 ± 3.45
7	Red soil : Coco peat (1:1)	35.0 ± 4.33

*Note: Readings were taken 5 weeks after planting. Experiment had 20 replicates and experiment was repeated at least thrice.*



**Figure 1:** Somatic embryogenesis in *Solanum virginianum* L. a. Embryogenic callus, b. Induction of polar tip of embryo, c. Root development from the polar tip of embryo, d. Shoot initiation from the opposite pole of cell, e. Embryo growing into a well-developed plantlet, f. Heart shaped somatic embryo, g. Torpedo stage of embryogenesis, h., i. and j. Developing shoot tips, k. Distinct shoot tip dissected out from callus mass.

present study revealed that different concentrations of BAP with IAA, IBA induced embryogenic callus followed by somatic embryogenesis. 2,4-D and NAA alone induced embryogenic callus which when subcultured on hormone free MS gave somatic embryogenesis. The protocol developed is efficient and can be exploited for large scale multiplication of this rare and endangered medicinal plant.

## References

- Rita P and Animesh DK. An updated overview on *Solanum xanthocarpum* Schrad and Wendl. International Journal of Research in Ayurveda and Pharmacy, 2 (2011): 730-735.
- Sharma N, Sharma AK and Zafar R. Kantikari: A prickly medicinal weed ~ Ecosensorium. Journal of Phytological Research, 9(2010): 13-17.
- Rane MH, Sahu NK, Ajoankar SS, Teli NC and Verma DR 2014. A Holistic Approach on Review of *Solanum virginianum* L. Research and Reviews:Journal of Pharmacy and Pharmaceutical Science (2014): 2322-0112.
- Okram Mukherjee Singh and Thokchom Prasanta Singh. 2010. Phytochemistry of *Solanum xanthocarpum*: an amazing traditional healer. *Journal of Scientific & Industrial Research*. 69:732-740.
- Gnana Sundari S, Rekha S and Parvathi A. Phytochemical evaluation of three species of *Solanum*. International Journal of research Ayurveda and Pharmacy, 4 (2013): 420-425.
- Patel P, Patel M, Saralai M and Gandhi T. Antiurolithiatic Effects of *Solanum xanthocarpum* Fruit Extract on Ethylene-Glycol-Induced Nephrolithiasis in Rats. J Young Pharm. 4(2012):164-170.
- Patel VB, Rathod IS, Patel JM and Brahmabhatt MR. Aanti-urolithiatic and natriuretic activity of steroidal constituents of *Solanum xanthocarpum*. Der Pharma Chemica, 2(2010): 173-176.
- Tupkari SV, Saoji AN and Deshmukh VK. Phytochemical study of *Solanum virginianum*. *Planta Medica*, 22(1972); 184-187.
- Pratap Kumar Sahu, Sweta Priyadarshini Pradhan, P Sudhir Kumar. Isolation, elucidation, and structure-activity relationships of phytoalkaloids from Solanaceae. *In: Studies in Natural Products Chemistry*, Elsevier. 72 (2002): 371-389.
- Galanas IT, Webb DT and Rosario O. Steroidal production by callus and cell suspension cultures of *Solanum aviculare* Frost. *Journal of Natural Products*, 47 (1984): 73-79.
- Marck TK. *In vitro* culture and the production of Solasodine *In: Biotechnology in Agriculture, Forestry, Medicinal and Aromatic Plant*, vol VII. Springer-Verlag, Berlin Germany (1989).
- Singh M and Chaturvedi R. An Efficient Protocol for Cyclic Somatic Embryogenesis in Neem (*Azadirachta indica* A. Juss) Proceedings of International Conference on Energy and Environment (2009) pp. 656-658.
- Krayer O and Briggs LH. Studies on *Solanum* alkaloids; II The anti-accelerator cardiac action of solasodine and some of its derivatives. *British Journal of Pharmacology*, (1950): 5- 517.
- Yadav SK, Kachhwaha S and Kothari SL 2010. Comparison of *in vitro* regeneration efficiency of leaf explants in response to different cytokinins and assessment of genetic uniformity of regenerated plants of *Solanum surattense* Burm.f. *African journal of Biotechnology*, 9(2010): 8991-8997.
- Vasil IK 2008. A history of plant biotechnology: from the cell theory of Schleiden and Schwann to biotech crops. *Plant cell Reports*, 27 (2008) :1423-40
- Rao KS 1996. Embryogenesis in flowering plants: recent approaches and prospects. *Journal of Biosciences*, 21 (1996): 827-841.
- Jimenez VM. Involvement of plant hormones and plant growth regulators on *in vitro* somatic embryogenesis. *I*, 47 (2005): 91-110.
- Ramaswamy N, Ugandhar T, Praveen M, Lakshman A, Rambabu M, Venkataiah P. *In vitro* propagation of medicinally important *Solanum surattense*. *Phytomorphology*, 54 (2004): 281-289.
- Pawar PK, Pawar CS, Narkhede BA, Teli NP, Bhalsing SR and Maheshwari VL. A technique for rapid micropropagation of *Solanum surattense* Burm. F. *Indian Journal of Biotechnology*, 1 (2002): 201-204.
- Ramaswamy N, Ugandhar T, Praveen M, Venkataiah P, Rambabu M, Upendar M and Subhash K. Somatic embryogenesis and plantlet regeneration from cotyledon and leaf explants of *Solanum surattense*. *Indian Journal of Biotechnology*, 4 (2005): 414-418.
- Prasad, R.N., Sharma, M., Sharma, A.K. and Chaturvedi, H.C. 1998. Androgenic stable somaclonal variant of *Solanum surttense* Burm. F. *Indian Journal of Experimental Biology*, 36: 1007-1012.
- Khan TI and Frost S. Floral biodiversity: a question of survival in the Indian thar desert. *The Environmentalist*, 21 (2001): 231-236.
- Murashige T and Skoog F. A revised medium for rapid

- growth and bioassays with tobacco tissue cultures. *Physiologia Plantarum*, 15 (1962): 473-497.
24. Sundar AN and Jawahar M 2011. *In vitro* plant regeneration from leaf and stem explants of *Solanum xanthocarpum* Schrad & Wendl. – An important medicinal herb. *Journal of Agricultural Technology*, 7(2011):301-306.
  25. Ramar K and Nandagopalan V. Rapid *in-vitro* propagation of medicinally important plant *Solanum surattense* Burm F. *International Journal of Pharmaceutical and Life Science*, 2 (2011): 499-501.
  26. Dantu PK, Tomar UK and Tripathi G. Somatic embryogenesis. *In: Cellular and Biochemical Science*. New Delhi: IK International House Pvt Ltd (2010): 892–908.
  27. Sharma P and Rajam MV 1995. Genotype, explants and position effect on organogenesis and somatic embryogenesis in egg plant (*Solanum melongena* L.). *Journal of Experimental Botany*, 46 (1995): 135-141.
  28. Ghatge SR, Kedage VV and Dixit GB 2008. Somatic embryogenesis and plant regeneration of a rare and multipurpose medicinal plant *HEMIDESMUS INDICUS* (L.) SCHULT. *Plant Cell Biotechnology and Molecular Biology*, 9 (2008): 71-74.
  29. Zimmerman JL. Somatic embryogenesis: A model for early development in higher plants. *Journal of Plant Cell*, 5 (1993): 141-143.
  30. Sharma SK and Millam S 2004. Somatic embryogenesis in *Solanum tuberosum* L.: A histological examination of key developmental stages. *Plant Cell Reports*, 23 (2004): 115–119.
  31. Nuray C, Yesim YM, Selay E and Ceren U 2009. Effects of different combinations and concentrations of growth regulators and photoperiod on somatic embryogenesis of *cucumis melo* var. *flexuosus*. *African Journal of Biotechnology*, 8 (2009): 6228-6232.